

FORM PTO-1390 (Modified)
(REV 11-98)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES

POELSTRA

DESIGNATED/ELECTED OFFICE (DO/EO/US)

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR 1.5)

CONCERNING A FILING UNDER 35 U.S.C. 371

Unknown 09/868585

INTERNATIONAL APPLICATION NO
PCT/NL98/00722INTERNATIONAL FILING DATE
21 December 1998PRIORITY DATE CLAIMED
21 December 1998

TITLE OF INVENTION

DIAGNOSIS OF SEPSIS USING THE LPS-BINDING MOIETY OF ALKALINE PHOSPHATASE

APPLICANT(S) FOR DO/EO/US

Klaas POELSTRA

Arie Jacob MAN IN 'T VELD

Dirk Klaas Fokke MEIJER

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371 (c) (2))
 - a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
- ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
- ☒ A copy of the International Search Report (PCT/ISA/210).
- ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired
 - d. ☒ have not been made and will not be made.
- ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3))
- ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4))
- ☒ A copy of the International Preliminary Examination Report (PCT/IPEA/409).
- ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).

Items 13 to 20 below concern document(s) or information included:

13. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
14. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
15. ☒ A **FIRST** preliminary amendment.
16. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
17. ☐ A substitute specification.
18. ☐ A change of power of attorney and/or address letter.
19. ☐ Certificate of Mailing by Express Mail
20. ☐ Other items or information:

Express Mail Label No. EL695230229US


U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR 1.5) 09/868585 Unknown		INTERNATIONAL APPLICATION NO PCT/NL98/00722		ATTORNEY'S DOCKET NUMBER POELSTRA	
--	--	---	--	---	--

21. The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) :				CALCULATIONS PTO USE ONLY	
<input type="checkbox"/> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$970.00 <input checked="" type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$840.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$690.00 <input type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$670.00 <input type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) \$96.00				ENTER APPROPRIATE BASIC FEE AMOUNT = \$860.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (e)).				\$130.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	36 - 20 =	16	x \$18.00	\$288.00	
Independent claims	8 - 3 =	5	x \$80.00	\$400.00	
Multiple Dependent Claims (check if applicable).				\$0.00	
TOTAL OF ABOVE CALCULATIONS				= \$1,678.00	
Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28) (check if applicable).				\$0.00	
SUBTOTAL				= \$1,678.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (f)).				\$0.00	
TOTAL NATIONAL FEE				= \$1,678.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable).				\$0.00	
TOTAL FEES ENCLOSED				= \$1,678.00	
				Amount to be:	\$
				refunded	
				charged	\$

<input checked="" type="checkbox"/>	A check in the amount of \$1,678.00	to cover the above fees is enclosed.
<input type="checkbox"/>	Please charge my Deposit Account No. _____ in the amount of _____	to cover the above fees.
A duplicate copy of this sheet is enclosed.		
<input checked="" type="checkbox"/>	The Commissioner is hereby authorized to charge any fees which may be required, or credit any overpayment to Deposit Account No. 080570 A duplicate copy of this sheet is enclosed.	

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:


 SIGNATURE

Anthony H. Handal/ Roger Pitt
 NAME

26,275/46/996
 REGISTRATION NUMBER

20 June 2001
 DATE

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

-----X
In re application of : Attorney Docket
No.: POELSTRA
Applicant: Klaas POELSTRA et al. :
Serial No.: (Unknown) U.S. National Phase :
of PCT/NL98/00722 GAU: Unknown
:
Filed: 21 December 1998 Examiner: Unknown
:
For: "DIAGNOSIS OF SEPSIS USING
THE LPS-BINDING MOIETY OF :
ALKALINE PHOSPHATASE"
:
June 20, 2001

-----X
BOX PCT
Assistant Commissioner for Patents
Washington, D.C. 20231

PRELIMINARY AMENDMENT

SIR:

Prior to calculation of the filing fee, please make the following
amendments, referring to the claims of the international application as amended
by the response to the written opinion dated January 19, 2001.

IN THE CLAIMS:

Please AMEND claims 1, 3-11, 14-15, 17, 19-24, 30-31 and 36 to read as follows:

1. (amended) A method of diagnosis of onset of endotoxemia or sepsis due to Gram negative bacterial infection said method comprising monitoring of the degree of AP occupancy of LPS (Lipopolysaccharide) binding sites on alkaline phosphatase in a sample of tissue or fluid derived from a patient, wherein the degree of AP (Alkaline Phosphatase) occupancy is associated with presence or absence of Gram negative bacterial infection.

3. (amended) A method according to claim 1, wherein the degree of AP occupancy of LPS binding sites on alkaline phosphatase in a sample or tissue or fluid derived from a patient, is monitored over a period of time, wherein a decline of the degree of AP occupancy indicates Gram negative bacterial infection.

4. (amended) A method according to claim 1, wherein the degree of AP occupancy of LPS binding sites on alkaline phosphatase in the sample is determined and wherein onset of decline in the degree of AP occupancy indicates onset of Gram negative bacterial infection.

5. (amended) A method according to claim 1 wherein the degree of AP occupancy of LPS binding sites on alkaline phosphatase may also indicate a

mixed or single infection of Gram negative and Gram positive bacteria.

6. (amended) A method according to claim 1 wherein the sample is subjected to binding with a ligand for the LPS binding site on alkaline phosphatase followed by a determination of the degree of binding of the ligand.

7. (amended) A method according to claim 1 wherein the ligand for the LPS binding site on alkaline phosphatase is selected from the group consisting of naturally occurring ligands, chemically modified or genetically modified derivatives of natural LPS binding site binding substances, chemically produced ligands.

8. (amended) A method according to claim 1, wherein the sample is subjected to binding with a ligand for the LPS binding site on alkaline phosphatase selected from LPS, Lipid A, and LPS binding site antibody against alkaline phosphatase, a Fab fragment with LPS binding site binding ability on alkaline phosphatase, a single chain fragment of an immunoglobulin having LPS binding site binding activity on alkaline phosphatase.

9. (amended) A method according to claim 1, wherein the LPS binding site binding ligand has at least the affinity for the LPS binding site of alkaline

phosphatase of LPS

10. (amended) A method according to claim 1 wherein the LPS binding site binding ligand has at least the affinity for the LPS binding site of alkaline phosphatase of lipid A.

11. (amended) A method according to claim 1 wherein the degree of AP occupancy of LPS binding sites on alkaline phosphatase is determined by assessment of the dephosphorylating capacity of alkaline phosphatase in the sample.

14. (amended) A method according to claim 1 wherein the sample is from a cholestasis free patient.

15. (amended) A method according to claim 1 wherein the method also comprises a further assay of a sample from the patient for another disease related to increase of alkaline phosphatase activity, said further assay employing a method avoiding determination of alkaline phosphatase level.

17. (amended) A method according to claim 1, wherein the sample is taken from an individual at risk of Gram negative bacterial infection.

19. (amended) A method according to claim 1 wherein the sample is taken from an individual during hospitalization.

20. (amended) A method according to claim 1 wherein the sample is taken a number of times over a period of time and the data are compared thus revealing the level of AP occupancy over time.

21. (amended) A method according to claim 1 wherein the period of time is as long as the individual is at risk of infection i.e. during hospitalization or post trauma recovery.

22. (amended) A method according to claim 1 wherein the result of the assay is compared to a standard value thus revealing whether the degree of AP occupancy is indicative of endotoxemia or sepsis or the risk thereof.

23. (amended) A method according to claim 1 wherein the sample is a sample selected from the group consisting of blood and tissue, said blood sample for example being serum, and the tissue being other than bone and said tissue for example being selected from liver and intestine.

24. (amended) A kit comprising alkaline phosphatase LPS binding site binding

ligand and instructions for carrying out an assay according to claim 1 and any additional component required for such assay being selected from the following group of said group consisting of detectable marker, buffer, containers, comparative samples, data charts e.g. standard curves and data concerning relevant data of alkaline phosphatase values.

30. (amended) Use of a compound with an LPS binding site of alkaline phosphatase such as LPS, lipid A or a ligand which binds to the LPS binding site of AP as described in claim 1 as active compound in a method of preparation for medicament for therapy or diagnosis.

31. (amended) Use of a compound with an LPS binding site of alkaline phosphatase such as LPS, lipid A or a ligand which binds to the LPS binding site of AP as described in claim 1 as active compound in a method of preparation for as medicament for therapy or diagnosis of endotoxemia or sepsis.


36. (amended) A method or use according to claim 1 wherein the method is carried out on a sample derived from the group of individuals consisting of a patient, an individual at risk of Gram negative bacterial infection, an individual prior to or after trauma, an individual during hospitalization.

REMARKS

The claim amendments are made in order to remove multiple dependencies.

Respectfully submitted,

By



Anthony H. Handal

Reg. No. 26,275

Roger Pitt

Reg. No. 46,996

Express Mail Label No. EL695230229US

"VERSION WITH MARKINGS TO SHOW CHANGES MADE"

1. (amended) A method of diagnosis of onset of endotoxemia or sepsis due to Gram negative bacterial infection said method comprising monitoring of the degree of AP occupancy of LPS (Lipopolysaccharide) binding sites on alkaline phosphatase in a sample of tissue or fluid derived from a patient, wherein the degree of AP (Alkaline Phosphatase) occupancy is associated with presence or absence of Gram negative bacterial infection.

3. (amended) A method according to claim 1 ~~or 2~~, wherein the degree of AP occupancy of LPS binding sites on alkaline phosphatase in a sample or tissue or fluid derived from a patient, is monitored over a period of time, wherein a decline of the degree of AP occupancy indicates Gram negative bacterial infection.

4. (amended) A method according to ~~any of the preceding claims~~ claim 1, wherein the degree of AP occupancy of LPS binding sites on alkaline phosphatase in the sample is determined and wherein onset of decline in the degree of AP occupancy indicates onset of Gram negative bacterial infection.

5. (amended) A method according to ~~any of the preceding claims~~ claim 1 wherein the degree of AP occupancy of LPS binding sites on alkaline phosphatase may also indicate a mixed or single infection of Gram negative and Gram positive bacteria.

6. (amended) A method according to ~~any of the preceding claims~~ claim 1 wherein the sample is subjected to binding with a ligand for the LPS binding site on alkaline phosphatase followed by a determination of the degree of binding of the ligand.

7. (amended) A method according to ~~any of the preceding claims~~ claim 1 wherein the ligand for the LPS binding site on alkaline phosphatase is selected from the group consisting of naturally occurring ligands, chemically modified or genetically modified derivatives of natural LPS binding site binding substances, chemically produced ligands.

8. (amended) A method according to ~~any of the preceding claims~~ claim 1, wherein the sample is subjected to binding with a ligand for the LPS binding site on alkaline phosphatase selected from LPS, Lipid A, and LPS binding site antibody against alkaline phosphatase, a Fab fragment with LPS binding site binding ability on alkaline phosphatase, a single chain fragment of an

immunoglobulin having LPS binding site binding activity on alkaline phosphatase.

9. (amended) A method according to ~~any of the preceding claims~~ claim 1, wherein the LPS binding site binding ligand has at least the affinity for the LPS binding site of alkaline phosphatase of LPS

10. (amended) A method according to ~~any of the preceding claims~~ claim 1 wherein the LPS binding site binding ligand has at least the affinity for the LPS binding site of alkaline phosphatase of lipid A.

11. (amended) A method according to ~~any of claims 1-5~~ claim 1 wherein the degree of AP occupancy of LPS binding sites on alkaline phosphatase is determined by assessment of the dephosphorylating capacity of alkaline phosphatase in the sample.

14. (amended) A method according to ~~any of the preceding claims~~ claim 1 wherein the sample is from a cholestasis free patient.

15. (amended) A method according to ~~any of claims 1-13~~ claim 1 wherein the method also comprises a further assay of a sample from the patient for another

disease related to increase of alkaline phosphatase activity, said further assay employing a method avoiding determination of alkaline phosphatase level.

17. (amended) A method according to ~~any of claims 1-13~~ claim 1, wherein the sample is taken from an individual at risk of Gram negative bacterial infection.

19. (amended) A method according to ~~any of the preceding claims~~ claim 1 wherein the sample is taken from an individual during hospitalization.

20. (amended) A method according to ~~any of the preceding claims~~ claim 1 wherein the sample is taken a number of times over a period of time and the data are compared thus revealing the level of AP occupancy over time.

21. (amended) A method according to ~~any of the preceding claims~~ claim 1 wherein the period of time is as long as the individual is at risk of infection i.e. during hospitalization or post trauma recovery.

22. (amended) A method according to ~~any of the preceding claims~~ claim 1 wherein the result of the assay is compared to a standard value thus revealing whether the degree of AP occupancy is indicative of endotoxemia or sepsis or the risk thereof.

23. (amended) A method according to ~~any of the preceding claims~~ claim 1 wherein the sample is a sample selected from the group consisting of blood and tissue, said blood sample for example being serum, and the tissue being other than bone and said tissue for example being selected from liver and intestine.

24. (amended) A kit comprising alkaline phosphatase LPS binding site binding ligand and instructions for carrying out an assay according to ~~any of the preceding claims~~ claim 1 and any additional component required for such assay being selected from the following group of said group consisting of detectable marker, buffer, containers, comparative samples, data charts e.g. standard curves and data concerning relevant data of alkaline phosphatase values.

30. (amended) Use of a compound with an LPS binding site of alkaline phosphatase such as LPS, lipid A or a ligand which binds to the LPS binding site of AP as described in ~~any of the preceding claims~~ claim 1 as active compound in a method of preparation for medicament for therapy or diagnosis.

31. (amended) Use of a compound with an LPS binding site of alkaline phosphatase such as LPS, lipid A or a ligand which binds to the LPS binding site of AP as described in ~~any of the preceding claims~~ claim 1 as active compound in

a method of preparation for as medicament for therapy or diagnosis of endotoxemia or sepsis.

36. (amended) A method or use according to ~~any of the preceding claims~~ claim 1 wherein the method is carried out on a sample derived from the group of individuals consisting of a patient, an individual at risk of Gram negative bacterial infection, an individual prior to or after trauma, an individual during hospitalization.

THE "SOCIETY"

11/PR'75

09/868585

JC18 Rec'd PCT/PTO 20 JUN 2001

PCT/NL98/00722

WO 00/37943

1

DIAGNOSIS OF SEPSIS USING THE LPS-BINDING MOIETY OF ALKALINE PHOSPHATASE

5 PRESENT STATE OF THE ART

Alkaline phosphatase (AP) is an ecto-enzyme bound via a glycosyl-phosphatidyl-inositol (GPI)-anchor to the plasma membrane of many cell types including endothelial and epithelial cells, osteoblasts, fibroblasts, and hepatocytes. The enzyme is also present in specific granula of neutrophils. Consequently, all organs
10 contain abundant alkaline phosphatase activity. In addition, enzyme activity is also present in plasma.

Increased levels of AP activity in blood are found when bone formation is evident for instance during embryogenesis and childhood (3). Also hepatic diseases like cholestasis are characterized by high serum AP levels and an enhanced expression of AP
15 activity upon plasma membranes of hepatocytes. This enzyme is therefore a generally accepted serum marker for the diagnosis of this disease (4). Release from neutrophilic granula after activation of this cell type may also contribute to the enhanced serum AP activity found during inflammatory processes. The physiological significance of this enzyme induction in liver diseases and inflammatory processes is unknown. A role of AP
20 during the process of bone formation has been documented in vitro and in vivo.

Recent studies have demonstrated that AP is able to dephosphorylate endotoxin or lipopolysaccharide (LPS). LPS is a constituent of the cell wall of Gram-negative bacteria and it is very toxic to man and most animal species. It induces fulminant inflammatory reactions which may be life threatening when these responses
25 occur within the bloodstream. This disease is referred to as 'sepsis' or 'systemic inflammatory response syndrome (SIRS)' and is lethal in approximately 35% of all cases. LPS may enter the bloodstream after bacterial infections, during hepatic diseases or when intestinal permeability is enhanced for instance during or after ischemic occurrences. High concentrations of LPS are present in the gastro-intestinal tract due to
30 production by E. coli bacteria, Enterobacteriaceae, and many other bacterial species. Since virtually no LPS is found in plasma, the gastrointestinal tract obviously has an effective barrier. This barrier is characterized by a high AP expression similar to the blood brain barrier where neuro endothelial cells also express a high AP activity.

An enhanced intestinal permeability for LPS has been documented in patients with gastro-intestinal perforations, appendicitis or peritonitis but also in patients with alcoholic abuse, after traumatic events such as burns, major surgical operations to the gastro-intestinal tract and after sepsis. Most of these latter cases are characterized by post-ischemic reperfusion injury to the intestinal wall and concomitant release of an array of inflammatory mediators which affect the vascular permeability in the intestine thus causing leakage of LPS from the intestinal lumen into the vena porta. In normal conditions the liver plays an important role in the removal of LPS from the circulation, in particular the Kupffer cell is important in this respect. However, in many hepatic diseases this function is impaired which in turn may lead to endotoxemia.

Phosphate groups within the lipid A moiety of LPS determine several biological responses to this bacterial product. The lipid A moiety is the relatively constant part of this otherwise highly variable molecule. Generally, two phosphate groups are present upon this lipid A moiety in addition to a variety of phosphate groups within the polysaccharide tail. Whereas phosphate groups within the polysaccharide tail do not influence the inflammatory response upon LPS, the phosphate groups within the lipid A moiety are of major importance. It has been demonstrated that mono-phosphoryl lipid A is a weak inducer of macrophage responses and is non-toxic to chickens and rabbits, whereas native, di-phosphoryl lipid A is a potent activator of macrophages and lethal to most mammals.

The dephosphorylation of LPS by alkaline phosphatase may therefore be important in vivo and this creates new insights into the physiological role of this enzyme. This new insight may have many therapeutic implications. Document PCT/NL94/00189 describes the therapeutic application of alkaline phosphatase to treat all kinds of the clinical complications induced by LPS.

DESCRIPTION OF THE INVENTION

In summary, the examples show that serum AP activity declines in patients with established Gram-negative sepsis, whereas serum AP activity is unaffected or even is slightly enhanced in patients with established Gram-positive sepsis. This new finding may be explained by the binding of LPS to AP enzymes as demonstrated herein. This binding either may cause a change in the distribution of AP, thus reducing half-life in

serum or it may affect the enzyme activity directly. Based on all these observations, and considering the new insight on the role of AP in vivo as an LPS detoxifying enzyme (described in document PCT/NL94/00189), it can be concluded that AP has a binding site for LPS and that the binding of this ligand to the enzyme has important implications.

5 Early detection of endotoxemia in patients with liver failure, severe burns or other trauma's (e.g. after organ transplantations) is important but rather complicated. Reliable and sensitive methods which measure free LPS in serum and which take the endotoxin detoxifying capacity of serum also into account are not available yet. In addition, methods determining the amount of LPS in serum do not give any information
10 about the preceding situation. This is particularly relevant in view of the short half-life of LPS in the bloodstream. Detection methods to measure the binding of LPS to AP as such are feasible and these methods can be applied for diagnostic purposes to discriminate between Gram-positive and Gram-negative sepsis or to assess the severity of the disease. The degree of occupation of the binding site appears to be an early marker for this disease. Thus a valuable new diagnostic tool is described here. In addition,
15 based upon the insight that AP has a specific binding domain for LPS, new LPS binding systems can be developed that bind to this domain. These domains are able to remove LPS from fluid or tissue. ligands interacting with AP may also be used for the purification of AP from crude or partially purified biological sources (e.g. milk, culture media, tissue samples) Also systems applying the LPS binding domain of AP can also be
20 used for therapeutic purposes.

Thus the invention describes the use of ligands to the LPS-binding site of alkaline phosphatase. Examples of such ligands are mono- or polyclonal antibodies against this ligand either in their natural occurring form or as Fab-fragments, single
25 chains, or otherwise modified immunoglobulins. In addition, labeled LPS, lipid A, derivatives of both substances or products mimicking the tertiary structure of lipid A can be used as ligands for the LPS binding site in AP enzymes. It is possible to design such on the basis of computer modelling. The product can be produced synthetically using chemical means. There are references detailing production of lipid A like structures using chemical means (1,2). Use of recombinant DNA technology to engineer the
30 required structure is also possible as is chemical modification of lipid A like structures. Measurement of the concentration of free LPS-binding domains on alkaline phosphatase for diagnostic purposes can be done using these products. Subsequently, this

concentration can be related to the total concentration of AP enzymes, or to the total enzyme activity in biological fluids which will yield additional information about the presence of LPS and the LPS-detoxifying capacity in biological fluids.

Thus the method according to the invention consists of diagnosis of onset of
5 endotoxemia or sepsis due to Gram negative bacterial infection said method comprising
monitoring of the degree of AP occupancy of LPS binding sites on alkaline phosphatase
in a sample of tissue or fluid derived from a patient, wherein the degree of AP
occupancy is associated with presence or absence of Gram negative bacterial infection.
Specifically such a method wherein the degree of AP occupancy of LPS binding sites
10 on alkaline phosphatase in the sample is lower than that of an equivalent sample type of
an individual free of Gram negative infection. By way of example the degree of AP
occupancy of LPS binding sites on alkaline phosphatase in a sample or tissue or fluid
derived from a patient, is monitored over a period of time, wherein a decline of the
degree of AP occupancy indicates Gram negative bacterial infection. In such a case the
15 degree of AP occupancy of LPS binding sites on alkaline phosphatase in the sample is
determined and when onset of decline in the degree of AP occupancy is ascertained this
indicates onset of Gram negative bacterial infection. It is also possible depending on the
values determined that the degree of AP occupancy of LPS binding sites on alkaline
phosphatase may also indicate a mixed infection of Gram negative and Gram positive
20 bacteria. The test may indicate the onset of sepsis but may also identify those individuals
at risk for sepsis e.g. those individuals at risk for sepsis e.g. persons with structural low
AP levels in tissue or blood, due to a genetic defect or due to a chronic illness such as
Rheumatoid arthritis, brain oedema, Alzheimer, ischemic heart diseases or inflammatory
bowel diseases. In addition the test may be used to assess (subclinical) infections in
25 pregnant women. These subclinical infections may provoke the HELLP syndrome or pre
eclampsia in the mother or affect the condition of the child by causing intraurine growth
retardation of the foetus or even inducing pre-term delivery. As a consequence the test
described here can be used to prevent pathology to the new born child. In a method
according to the invention the sample to be determined can be subjected to binding with
30 a ligand for the LPS binding site on alkaline phosphatase followed by determination of
the degree of binding of the ligand. Any ligand with the required specificity can be used.
The selection will depend on the type of sample to be tested and the degree of sensitivity
to be reached. Quite specifically suitable ligands can be selected LPS, Lipid A, an LPS

binding site antibody against alkaline phosphatase, a Fab fragment with LPS binding site binding ability on alkaline phosphatase, a single chain fragment of an immunoglobulin having LPS binding site binding activity on alkaline phosphatase. Alternatively other ligands e.g. developed on the basis of computer modelling of the binding system of LPS and lipid A with alkaline phosphatase can also be used. The ligands may be naturally occurring molecules or synthetic molecules. The synthetic molecules may be produced using chemical or biochemical i.e. recombinant DNA technology methods. Preferably a method according to the invention employs an LPS binding site binding ligand with at least the affinity for the LPS binding site of alkaline phosphatase of LPS. Also a method according to the invention may suitably employ an LPS binding site binding ligand with at least the affinity for the LPS binding site of alkaline phosphatase of lipid A. The required degree of specificity and affinity can be ascertained using standard tests for comparing specificity and affinity of a compound for a substrate. In a method according to the invention the degree of AP occupancy of LPS binding sites on alkaline phosphatase can be determined by assessment of the dephosphorylating capacity of alkaline phosphatase in the sample. In such a method the ratio of dephosphorylating alkaline phosphatase to non dephosphorylating alkaline phosphatase is determined. The ratio can be determined using the values obtained by assessment of total alkaline phosphatase activity using biochemical methods to determine dephosphorylating activity and by assessment of total amounts of alkaline phosphatase using antibodies and calculating the ratio of these values. There are numerous routes available to the skilled person to carry out such assessments. We refer by way of example to references 5 and 6. The content hereof is incorporated by reference. It will immediately be apparent what other alternative known methods can be applied to arrive at the required data.

The sample to be used in the method according to the invention can be a sample derived from a patient. Such a sample can be from blood or tissue. The sample must be a sample treated such that any alkaline phosphatase present is not removed prior to the assay. Such a sample can be selected from the group consisting of blood and tissue. The blood sample can for example be serum. The tissue sample is preferably other than bone as alkaline phosphatase levels are generally high in bone. The tissue that is suitable for example can be selected from liver and intestines. A biops can be taken in a manner known per se. Blood and serum samples may also be prepared in the usual manner.

The method according to the invention can be carried out with a sample from a cholestasis free patient. The method according to the invention can also be carried out on a patient with cholestasis. The values determined for the degree of LPS binding can be compared to standard values of healthy and/or sick patients to ascertain the correct diagnosis. Alternatively the method according to the invention can comprise carrying out the assay described above in any of the described embodiments and in addition also comprising a further assay of a sample from the patient for another disease related to increase of alkaline phosphatase activity such as cholestasis. The further assay must preferably then employ a method avoiding determination of alkaline phosphatase level. Numerous articles describing such alternative assays for alkaline phosphatasae associated diseases are available in articles covering said diseases and the content thereof is incorporated by reference. A suitable embodiment of the method according to the invention wherein the further assay is carried out is when the alkaline phosphatase assay according to the invention reveals no decline in AP occupancy of LPS binding sites of alkaline phosphatase. Suitably the method according to any of the embodiments of the invention comprises a method wherein the sample is taken from an individual at risk of Gram negative bacterial infection. The sample can for example be taken from an individual either both prior to and following trauma. It can also be taken shortly after having undergone trauma. Quite interesting cases of trauma concern surgery, burns and ischemic trauma. As hospitals form notorious sources of infection suitably the method according to the invention can comprise assessment of a sample taken from an individual during hospitalisation.

In a method according to the invention the sample can be taken a number of times over a period of time. The data can be compared with each other thus revealing the level of AP occupancy over time. The values can however also be compared with a standard value for a healthy or sick individual. The values can also be correlated to various characteristics of the patient e.g. gender, age, body weight, tissue or fluid type to be tested. The method thus reveals whether the degree of AP occupancy is indicative of endotoxemia or sepsis. A method according to the invention can comprise taking samples over a period of time as long as the individual is at risk of infection i.e. during hospitalisation or post trauma recovery.

The invention also envisages kits for carrying out the diagnostic method according to the invention. Such a kit can comprise alkaline phosphatase LPS binding

site binding ligand and instructions for carrying out an assay according to any of the abovementioned embodiments. The kit can further comprise any additional components required for such assay e.g. detectable marker, buffer, containers. Also suitable as additions to the kit are comparative samples or data charts e.g. standard curves or data concerning relevant data of alkaline phosphatase values. A kit comprising alkaline phosphatase LPS binding site binding ligand for carrying out an assay according to any of the abovementioned embodiments and any additional component required for such assay being selected from the following group consisting of detectable marker, buffer, containers, comparative samples, data charts e.g. standard curves or data concerning relevant data of alkaline phosphatase values is also part of the invention. The ligands envisaged have been described in the embodiments of the method as disclosed above and in the claims and examples.

In addition the invention covers a method for removing LPS from tissue or fluid said method comprising contacting the LPS binding site of alkaline phosphatase with the tissue or fluid to be treated followed by separation of the LPS binding site and the tissue or fluid after the the LPS binding site has bound the LPS present in the fluid or tissue, with the proviso the ligand is neither alkaline phosphatase nor a derivative of alkaline phosphatase having dephosphorylating activity. Thus the LPS can be removed from the tissue or fluid. Clearly the binding must occur under circumstances that do not damage the fluid or tissue to be purified. Such conditions will be readily apparent to the skilled person used to dealing with such tissue or fluid. By way of example the LPS-binding domain of AP itself can be coupled to a non-soluble carrier to remove LPS from tissue or fluids. Suitable fluids to be treated are cell-culture media, liquids used for organ perfusion or preservation, blood, milk or any other therapeutic product administered to a person. Ligands which bind to the LPS binding site of AP may also be used for purification of AP itself from tissue or body fluids or other biological production systems.

The invention also covers a method for therapy of endotoxemia or sepsis said method comprising administration of a pharmaceutically effective amount of the LPS binding site of alkaline phosphatase in a systemically acceptable form with the proviso the ligand is neither alkaline phosphatase nor a derivative of alkaline phosphatase having dephosphorylating activity. Any conventional medicinal dosage form will suffice. The dosage regime to be followed will depend on the condition of the patient, gender,

age, body weight by way of example and the physician will be able to determine the best regime to apply. The LPS-binding domain as such or in the form of a derivative of alkaline phosphatase, said derivative not having dephosphorylating activity may be applied therapeutically to treat a patient with a systemic inflammatory response syndrome or to prevent endotoxemia in patients at risk for such a complication. Binding of LPS in blood to this particular domain may prevent the interaction of LPS with other endogenous receptors in serum and upon plasma membranes of endothelial cells or macrophages such as LPS-binding protein, CD-14 and scavenger receptors. As a consequence, the inflammatory response upon LPS will be attenuated.

The invention also covers other therapy and diagnostic applications of the ligands described in the description of the invention and a method of purification of AP itself using LPS or any AP binding derivative thereof i.e. the other ligands mentioned.

EXPERIMENTAL DATA

EXPERIMENT I

These reports prompted us to examine the serum levels of AP during endotoxemia. It is generally accepted that AP is a marker of liver damage, and LPS is known to induce directly or indirectly damage to this organ. An increase in serum AP activity in patients with sepsis has been frequently documented and was also expected in mice with endotoxemia. However, after intravenous administration of LPS from E.coli (strain 055:B5, Sigma Chemical Co, USA, 2.5 mg/kg b.w.) to Balb/c mice (male, 20-25 g) at t=0 a rapid and sharp decline in serum AP activity was observed as compared to mice receiving saline (fig. 1). A similar decline was found after intraperitoneal administration of E.coli bacteria (ATCC 25922, 2×10^9 CFU) to mice (fig. 1).

Evaluation of clinical data of patients with recurrent endotoxemia revealed also a rapid primary sharp decline in serum AP activity, followed by a strong increase in the following days. It is believed that the first sharp decline of AP is caused by LPS. This decline enables residual LPS to evoke its inflammatory action thereby activating the destructive responses seen in sepsis patients. Clinically this primary reaction of AP decline has hitherto not been observed. Therefore to enable a fast detection of sepsis fast detection of declining AP levels is of most importance. The method described in the present document enables the measurement of such a fast kinetic change in AP levels. The basis in this lies in the fact that free and thereby fully active AP in plasma declines rapidly upon binding to LPS. The secondary increase in serum AP levels seen in most

patients may either reflect liver damage or may be induced by interleukin 6 or other inflammatory mediators which attenuate the expression of AP activity as recent studies have pointed out.

EXPERIMENT II

We examined subsequently 64 patients in the intensive care unit suffering from sepsis. The pathogenic bacteria causing the sepsis syndrome were identified in all of these cases. Patients with endotoxemia (n=16) were compared with patients with Gram-positive bacterial infections (n=30), which lack LPS in serum. The patients had a variety of underlying diseases, but patients with cholestasis, which is associated with an increased AP excretion, were excluded. Blood samples were routinely assayed for AP during hospitalization. Changes in serum AP levels were related to day 0, that is, the day of admission to the intensive care unit of the hospital. We calculated the change in serum AP activity, relative to day 0, before and after the blood samples became positive for bacteria or bacterial products (LPS). These studies demonstrate that a decline in serum AP activity is an early marker of sepsis: the decrease preceded the bacteremia (fig. 2). Moreover, the data show that Gram-negative bacterial infections can be discriminated from Gram-positive bacterial infections because patients with established Gram-negative infections displayed a decline in serum AP activity (mean slope= -21.5) whereas patients with a proven Gram-positive bacterial sepsis showed a slight increase (mean slope= 10.9) relative to day zero. Note that at day zero (admittance to Intensive Care) the actual value of serum AP may have been affected already. Patients with a mixed infection (both Gram-positive and Gram-negative, n= 18) showed an intermediate change in the serum AP levels (mean slope= 0.6). The rapid diagnosis of Gram-positive versus a Gram-negative bacterial infection is particularly important in acute situations where an immediate start an effective antibacterial therapy is essential. To date, this diagnosis can only be made after a successful culture of the pathogenic bacteria, which takes usually two days.

EXPERIMENT III

We hypothesized that the decline in serum AP levels was caused by an increased turn-over of the enzyme. We therefore examined whether a direct binding between LPS and AP enzymes occurred and whether this binding would influence the

enzyme activity. Placental AP (6.7 U/ml), isolated from fresh human placenta's was incubated with LPS from E.coli (strain 055:B5, Sigma Chemical co; 1 mg /ml) for 10 minutes at room temperature. Subsequently, AP was subjected to FPLC analysis. The FPLC was equipped with an anion-exchange column (mono-Q HR 5/5, Pharmacia, Sweden) and UV detection (280 nm). PlAP was injected in 0.2 M Tris/HCl buffer (pH 7.4) containing MgCl₂ with or without LPS. Elution was performed in Tris/HCl buffer (pH 7.4) with a flow rate of 0.5 ml/min and a NaCl gradient ranging from 0.0 to 0.45 M in 30 minutes. Retention time was measured. The retention time of plAP samples containing no LPS was 20 minutes, whereas the retention time shifted to 18 minutes after co-incubation with LPS. The shorter retention time of plAP samples containing LPS reflects an decreased negative charge of the complex, which indicates binding of LPS to AP enzymes thus neutralizing charges within the proteins.

EXPERIMENT IV

To assess whether product inhibition occurred, which might explain the decline in serum AP activity, we tested the effect of unphosphorylated lipid A, which is the end-product of the enzyme reaction, upon AP activity. Formalin/macrodex-fixed cryostat sections (4 um) of the rat intestine, kidney and placenta were incubated with LPS, MgCl₂ and lead-nitrate (Pb₂NO₃) in 0.2 M Tris/maleic acid buffer (pH 7.4) to demonstrate phosphatase activity according to standard procedures (K. Poelstra et al. Lab. Invest 76:319, 1997, Am J. Pathol. 151: 1163, 1997).

As documented before, rat intestinal and renal cryostat sections expressed abundant LPS dephosphorylating enzyme activity. LPS-dephosphorylating activity in the rat placenta was less abundant. The localization of this enzyme activity was similar to the distribution of AP activity in all three organs. However, pre-incubation of these cryostat sections with lipid A (Monophosphoryl lipid A from Salmonella minnesota R595, List Biological Laboratories Inc., Campbell, USA; 1 mg/ml) strongly attenuated intestinal and renal phosphatase activity when LPS was subsequently added as substrate. No phosphatase activity was detectable in placenta's pre-incubated with lipid A.

From these data it has become clear that AP upon binding to LPS is temporarily not available for dephosphorylation of additional LPS. This may be explained by relatively high affinity of LPS to the binding site of AP and vice versa.

LEGENDS TO THE FIGURES:

Figure 1:

Serum alkaline phosphatase levels in mice after administration of saline (i.v.), LPS (i.v.) or E.coli bacteria (i.p.) at $t=0$ (n=4 per group). The diamond signifies saline, the square signifies E coli, the triangle signifies LPS. t is set in hr along x axis and AP activity is represented in U/L on y-axis

Figure 2:

Changes in serum AP levels relative to the day of admission to the intensive care unit of the hospital, before and after the blood samples of patients became positive for bacteria or bacterial products (LPS). Patients were divided into two groups: patients with an established Gram-negative bacterial infection (n=16) and patients with an established Gram-positive bacterial infection (n=30). Patients with a mixed bacterial infection (n=18) or with SIRS without identification of the bacterium are not presented here. The dark symbol signifies Gram negative. The light symbol signifies Gram positive.

References:

Chemical derivatives of LPS of lipid A:

1. Qureshi N; Jarvis B; Takayama K; Sattar N; Hofman J; Stutz P: Natural and synthetic LPS and lipid a analogs or partial structures that antagonize or induce tolerance to LPS. *Prog Clin Biol Res* 1998; 397: 289-300

2. Brandenburg K; Kusumoto S; Seydel U: Conformational studies of synthetic lipid A analogues and partial structures by infrared spectroscopy. *Biochim Biophys Acta*. 1997; 1329; 183-201

Conditions of enhanced serum AP levels:

3. Ross PD; Knowlton W: Rapid bone loss is associated with increased levels of biochemical markers. *J Bone Miner Res*. 1998; 13; 297-302

4. Dourakis SP; Mayroyannis C; Alexopoulou A; Hadziyannis SJ: Prolonged cholestatic jaundice after endoscopic retrograde cholangiography. *Hepatogastroenterology* 1997; 44; 677-80

Methods to determine serum AP:

5. Peake MJ; Pejakovic M; White GH: Quantitative method for determining serum alkaline phosphatase isoenzyme activity: estimation of intestinal component. J Clin pathol 1988; 41; 202-206
- 5 6. McComb RB; Bowers GN; Posen S: Measurement of alkaline phosphatase activity. In: Alkaline Phosphatase, Plenum Press, New York and London, 1979: Ch 7, pp 289-372.

10322-505050

CLAIMS

1. A method of diagnosis of onset of endotoxemia or sepsis due to Gram negative bacterial infection said method comprising monitoring of the degree of AP occupancy of LPS binding sites on alkaline phosphatase in a sample of tissue or fluid derived from a patient, wherein the degree of AP occupancy is associated with presence or absence of Gram negative bacterial infection.

2. A method according to claim 1, wherein the degree of AP occupancy of LPS binding sites on alkaline phosphatase in the sample is lower than that of an equivalent sample type of an individual free of Gram negative infection.

3. A method according to claim 1 or 2, wherein the degree of AP occupancy of LPS binding sites on alkaline phosphatase in a sample or tissue or fluid derived from a patient, is monitored over a period of time, wherein a decline of the degree of AP occupancy indicates Gram negative bacterial infection.

4. A method according to any of the preceding claims, wherein the degree of AP occupancy of LPS binding sites on alkaline phosphatase in the sample is determined and wherein onset of decline in the degree of AP occupancy indicates onset of Gram negative bacterial infection.

5. A method according to any of the preceding claims wherein the degree of AP occupancy of LPS binding sites on alkaline phosphatase may also indicate a mixed or single infection of Gram negative and Gram positive bacteria.

6. A method according to any of the preceding claims wherein the sample is subjected to binding with a ligand for the LPS binding site on alkaline phosphatase followed by determination of the degree of binding of the ligand.

7. A method according to any of the preceding claims wherein the ligand for the LPS binding site on alkaline phosphatase is selected from the group consisting of naturally occurring ligands, chemically modified or genetically modified derivatives of natural LPS binding site binding substances, chemically produced ligands.

8. A method according to any of the preceding claims, wherein the sample is subjected to binding with a ligand for the LPS binding site on alkaline phosphatase selected from LPS, Lipid A, an LPS binding site antibody against alkaline phosphatase, a Fab fragment with LPS binding site binding ability on alkaline phosphatase, a single chain fragment of an immunoglobulin having LPS binding site binding activity on

alkaline phosphatase.

9. A method according to any of the preceding claims, wherein the LPS binding site binding ligand has at least the affinity for the LPS binding site of alkaline phosphatase of LPS.

5 10. A method according to any of the preceding claims wherein the LPS binding site binding ligand has at least the affinity for the LPS binding site of alkaline phosphatase of lipid A.

10 11. A method according to any of claims 1-5 wherein the degree of AP occupancy of LPS binding sites on alkaline phosphatase is determined by assessment of the dephosphorylating capacity of alkaline phosphatase in the sample.

12. A method according to claim 11, wherein the ratio of dephosphorylating alkaline phosphatase to non dephosphorylating alkaline phosphatase is determined.

13. A method according to claim 12, wherein the ratio is determined using the values obtained by assessment of total alkaline phosphatase activity using biochemical methods to determine dephosphorylating activity and by assessment of total amounts of alkaline phosphatase using e.g. antibodies or otherwise discriminating entities and calculating the ratio of these values.

14. A method according to any of the preceding claims wherein the sample is from a cholestasis free patient.

15. A method according to any of claims 1-13 wherein the method also comprises a further assay of a sample from the patient for another disease related to increase of alkaline phosphatase activity, said further assay employing a method avoiding determination of alkaline phosphatase level.

16. A method according to claim 15 wherein the further assay is carried out when no decline in AP occupancy of LPS binding sites of alkaline phosphatase according to the method of any of claims 1-14 is detected.

17. A method according to any of the preceding claims, wherein the sample is taken from an individual at risk of Gram negative bacterial infection.

18. A method according to claim 17 wherein the sample is taken from an individual either both prior to and following trauma or shortly after having undergone trauma, wherein the trauma in particular concerns surgery, burns or ischemic traumas.

19. A method according to any of the preceding claims wherein the sample is taken from an individual during hospitalisation.

20. A method according to any of the preceding claims wherein the sample is taken a number of times over a period of time and the data are compared thus revealing the level of AP occupancy over time.

21. A method according to any of the preceding claims wherein the period of time is as long as the individual is at risk of infection i.e. during hospitalisation or post trauma recovery.

22. A method according to any of the preceding claims wherein the result of the assay is compared to a standard value thus revealing whether the degree of AP occupancy is indicative of endotoxemia or sepsis or the risk thereof.

23. A method according to any of the preceding claims wherein the sample is a sample selected from the group consisting of blood and tissue, said blood sample for example being serum, said tissue being other than bone and said tissue for example being selected from liver and intestine.

24. A kit comprising alkaline phosphatase LPS binding site binding ligand and instructions for carrying out an assay according to any of the preceding claims and optionally any additional components required for such assay e.g. detectable marker, buffer, containers and comparative samples or data charts e.g. standard curves or data concerning relevant data of alkaline phosphatase values.

25. A kit comprising alkaline phosphatase LPS binding site binding ligand for carrying out an assay according to any of the claims 1-23 and any additional component required for such assay being selected from the following group consisting of detectable marker, buffer, containers, comparative samples, data charts e.g. standard curves or data concerning relevant data of alkaline phosphatase values.

25. A method for therapy of endotoxemia or sepsis said method comprising administration of a pharmaceutically effective amount of the LPS binding site of alkaline phosphatase in a systemically acceptable form with the proviso the ligand is neither alkaline phosphatase nor a derivative of alkaline phosphatase having dephosphorylating activity.

26. A method for removing LPS from tissue or fluid said method comprising contacting the LPS binding site of alkaline phosphatase with the tissue or fluid to be treated followed by separation of the LPS binding site and the tissue or fluid after the the LPS binding site has bound the LPS present in the fluid or tissue, with the proviso the ligand is neither alkaline phosphatase nor a derivative of alkaline phosphatase having

dephosphorylating activity.

27. A method of purification of AP itself from tissue or body fluids or other biological production systems with a compound with an LPS binding site of alkaline phosphatase such as LPS, lipid A or other ligand as described in the preceding claims.

28. Use of a compound with an LPS binding site of alkaline phosphatase such as LPS, lipid A or other ligand as described in the preceding claims as active compound in a medicament for therapy or diagnosis.

29. Use of a compound with an LPS binding site of alkaline phosphatase such as LPS, lipid A or other ligand as described in the preceding claims as active compound in therapy or diagnosis.

fig - 1





fig - 2





Paul Morofsky

District of Columbia Office
American Center - Tysons Corner
Suite 500
8300 Boone Boulevard
Vienna, Virginia 22182

HANDAL & MOROFSKY
Counsellors at Law
80 Washington Street
Norwalk, Connecticut 06854

Telephone: 203-838-8589
Facsimile: 203-838-8794
Telex: 429283 PATSLAW

JC05 R PCT/PTO 26 JUL 2001 #3

July 24, 2001

BOX MISSING PARTS

Assistant Commissioner for Patents
Washington, D.C. 20231

Re: Klaas POELSTRA et al.
U.S. Patent Application No.: 09/868,585
National Phase of International Application No. PCT/NL98/00722
Dated: December 21, 1998
"DIAGNOSIS OF SEPSIS USING THE LPS-BINDING MOIETY OF ALKALINE
PHOSPHATASE" Attorney Docket No. POELSTRA

SIR:

We enclose herewith:

- ☒ Combined Declaration and Power of Attorney;
- ☒ Acknowledgment Postcard.

CORRESPONDENCE INDICATION: Please recognize the following address as the
correspondence address:



00545

PATENT TRADEMARK OFFICE

The Commissioner is hereby authorized to charge payment of the fees associated
with this communication or credit any overpayment to Deposit Account No. 08-0570.

The applicants hereby petition under 37 CFR 1.136 or other applicable rule to have
the response period extended the number of months necessary to render the attached
communication timely if a petition is required.

Respectfully submitted,

Anthony H. Handal
Reg. No. 26,275
Roger Pitt
Reg. No. 46,996

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope, postage prepaid,
addressed to the Assistant Commissioner for Patents Washington, D.C. 20231 on July 24, 2001

Anthony H. Handal Reg. No. 26,275
Roger Pitt Reg. No. 46,996



COMBINED DECLARATION AND POWER OF ATTORNEY

ORIGINAL DESIGN, NATIONAL STAGE OF PCT OR CIP APPLICATION)

As a below named inventor, I hereby declare that

My residence, post office address and citizenship are as stated below next to my name, I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Diagnosis of sepsis using the LPS-binding moiety of alkaline phosphatase

the specification of which: (complete (a), (b) or (c) for type of application)

REGULAR OR DESIGN APPLICATION

a. ☐ is attached hereto.
b. ☐ was filed on _____ as Application
Serial No. _____ and was amended on _____
(if applicable)

PCT FILED APPLICATION ENTERING NATIONAL STAGE

c. ☒ was described and claimed in International application No. PCT/NL98/00722
filed on 21 December 1998
and as amended on _____ (if any)

ACKNOWLEDGEMENT OF REVIEW OF PAPERS AND DUTY OF CANDOR

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, paragraph 1.56(a).

In compliance with this duty there is attached an information
disclosure statement 37 CFR 1.97

PRIORITY CLAIM

I hereby claim foreign priority benefits under Title 35, United States Code paragraph 119 of any foreign application (s) for patent of inventor's certificate listed below and have also identified below any foreign application for patent of inventor's certificate having a filing date before that of the application on which priority is claimed.

(complete (d) or (e))

- d. ☐ no such applications have been filed
e. ☐ such applications have been filed as follows

**EARLIEST FOREIGN APPLICATION(S), IF ANY FILED WITHIN 12 MONTHS
(6 MONTHS FOR DESIGN) PRIOR TO SAID APPLICATION**

Country	Application Number	Date of filing (day, month, year)	Date of Issue (day, month, year)	Priority claimed

**ALL FOREIGN APPLICATION(S), IF ANY FILED MORE THAN 12 MONTHS
(6 MONTHS FOR DESIGN) PRIOR TO SAID APPLICATION**

CONTINUATION-IN-PART

(Complete this part only if this is a continuation-in-part application)

I hereby declare claim the benefit under Title 35, United States code, paragraph 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claim of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, paragraph 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, paragraph 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.) (Filing date) (Status) (patented, pending, abandoned)

(Application Serial No.) (Filing date) (Status) (patented, pending, abandoned)

POWER OF ATTORNEY

As a named inventor, I hereby appoint the following attorney and agent to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Anthony H. Handal, Reg. No. 26,275


Send Correspondence to: Handal & Morofsky, Counselors at Law,
80 Washington Street
NORWALK, Connecticut 06854

Direct telephone calls to: (203) 838-8589

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that wilful false statements and the like so made are punishable by fine or imprisonment, or both under Section 1001 of Title 18 of the United States Code and that such wilful false statements may jeopardize the validity of the application or any patent issued thereon.

1-00
Full name of sole or first inventor: POELSTRA, Klaas

Inventor's signature



Date


Country of Citizenship: the Netherlands

Residence: BUITENPOST, the Netherlands NLX

Post Office Address: Kruirad 2, NL-9285 MT BUITENPOST, the Netherlands

2-00
Full name of third inventor: MEIJER, Dirk Klaas Fokke

Inventor's signature



Date


Country of Citizenship: the Netherlands

Residence: Groningen, the Netherlands NLX

Post Office Address: Parklaan 17, NL-9724 AN GRONINGEN, the Netherlands

3-00
Full name of second inventor: MAN IN 'T VELD, Arie Jacob

Inventor's signature



Date

Country of Citizenship: The Netherlands

Residence: Rotterdam, the Netherlands NLX

Post Office Address: Laan van Ypenhof 93, NL-3062 ZN ROTTERDAM, the Netherlands

CHECK PROPER BOX(ES) FOR ANY ADDED PAGE(S) FORMING A PART OF THIS DECLARATION